

5-2016

# A fractional boundary value problem

Grant Yost

University of Tennessee at Chattanooga, [ddw497@mocs.utc.edu](mailto:ddw497@mocs.utc.edu)

Follow this and additional works at: <http://scholar.utc.edu/honors-theses>



Part of the [Mathematics Commons](#)

---

## Recommended Citation

Yost, Grant, "A fractional boundary value problem" (2016). *Honors Theses*.

This Theses is brought to you for free and open access by the Student Research, Creative Works, and Publications at UTC Scholar. It has been accepted for inclusion in Honors Theses by an authorized administrator of UTC Scholar. For more information, please contact [scholar@utc.edu](mailto:scholar@utc.edu).

A Fractional Boundary Value Problem

Grant Yost

Departmental Honors Thesis

The University of Tennessee at Chattanooga

Mathematics Department

Project Director: Dr. John Graef

Examination Date: 6 November, 2015

Members of Examination Committee:

  
\_\_\_\_\_

Project Director

  
\_\_\_\_\_

Department Examiner

  
\_\_\_\_\_

Department Examiner

  
\_\_\_\_\_

Liason, Departmental Honors Committee

  
\_\_\_\_\_

Chair, Departmental Honors Committee

We consider the boundary value problem consisting of the fractional differential equation

$$-D_{0+}^{\alpha}z + aD_{0+}^{\beta}z = w(t)f(t, z), \quad 0 < t < 1, \quad (1)$$

and the boundary conditions

$$z(0) = z'(0) = \dots = z^{(n-2)}(0) = 0, \quad z(1) = k, \quad (2)$$

where  $k \in \mathbb{R}$ ,  $a \in \mathbb{R}$ ,  $n \in \mathbb{N}$ ,  $0 \leq \beta \leq n - 2$ ,  $n - 1 < \alpha < n$ ,  $w \in L[0, 1]$  with  $w(t) \geq 0$ , and  $f \in C([0, 1] \times \mathbb{R}, \mathbb{R})$ . Here,  $D_{0+}^{\gamma}h$  is the  $\gamma$ -th left Riemann-Liouville fractional derivative of  $h : [0, 1] \rightarrow \mathbb{R}$  defined by

$$(D_{0+}^{\gamma}h)(t) = \frac{1}{\Gamma(l - \gamma)} \frac{d^l}{dt^l} \int_0^t (t - s)^{l-\gamma-1} h(s) ds, \quad l = \lfloor \gamma \rfloor + 1,$$

whenever the right-hand side exists with  $\Gamma(\cdot)$  being the Gamma function.

Fractional differential equation models such as this are relevant in various science and engineering fields. Problems like this have been solved by first finding the Green's function for  $z$  in similar equations such as

$$-D_{0+}^{\alpha}z = f(t, z).$$

The Green's function (denoted by  $G$ ) was developed as a method to solve boundary value problems with linear homogeneous equations (i.e. when  $f(t, z) = 0$ ). However, when the right side of equation is a function of  $t$ , say  $w(t)f(t, z(t))$ , the unique solution to the problem is given by

$$u(t) = \int_0^1 G(t, s)w(s)f(s, z(s))ds,$$

where  $G$  is the solution to the similar problem with the right hand side equal to zero.

We try to solve problem (1)-(2) by splitting it into the two following problems:

$$\begin{cases} -D_{0+}^{\alpha}u + aD_{0+}^{\beta}u = w(t)f(t, u), & 0 < t < 1, \\ u(0) = u'(0) = \dots = u^{(n-2)}(0) = 0, & u(1) = 0, \end{cases} \quad (3)$$

$$\begin{cases} -D_{0+}^{\alpha}v + aD_{0+}^{\beta}v = 0, & 0 < t < 1, \\ v(0) = v'(0) = \dots = v^{(n-2)}(0) = 0, & v(1) = k. \end{cases} \quad (4)$$

Note that if  $u(t)$  is a solution of (3) and  $v(t)$  is a solution of (4), then  $z(t) = u(t) + v(t)$  is a solution of (1)-(2).

The solution of (3) is found by using the Green's Function:  $G : [0, 1] \times [0, 1] \rightarrow \mathcal{R}$ , which was obtained in [3]:

$$G(t, s) = \begin{cases} \frac{[t(1-s)]^{\alpha-1} E_{\alpha-\beta, \alpha}[at^{\alpha-\beta}] E_{\alpha-\beta, \alpha}[a(1-s)^{\alpha-\beta}]}{E_{\alpha-\beta, \alpha}[a]}, & 0 \leq t \leq s \leq 1, \\ \frac{[t(1-s)]^{\alpha-1} E_{\alpha-\beta, \alpha}[at^{\alpha-\beta}] E_{\alpha-\beta, \alpha}[a(1-s)^{\alpha-\beta}]}{E_{\alpha-\beta, \alpha}[a]} \\ \quad - (t-s)^{\alpha-1} E_{\alpha-\beta, \alpha}[a(t-s)^{\alpha-\beta}], & 0 \leq s \leq t \leq 1, \end{cases} \quad (5)$$

with  $|a| < \Gamma(\alpha - \beta + 1)$ . Here, the Green's Function is given in terms of the known Mittag-Leffler function  $E_{\gamma, \delta} : \mathcal{C} \rightarrow \mathcal{C}$ :

$$E_{\gamma, \delta}[z] = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\gamma n + \delta)}.$$

Using this function, we find the solution to (3) to have the form:

$$u(t) = \int_0^1 G(t, s)w(s)f(s, u(s))ds$$

The solution to (4) is easily seen to be:

$$v(t) = \frac{kt^{\alpha-1}E_{\alpha-\beta,\alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta,\alpha}(a)}.$$

Note that by Lemma 4.3 of [3], the Mittag-Leffler function,  $E$ , is increasing, so  $|v(t)| \leq k$  for  $t \in [0, 1]$ .

The solution to (1)-(2) is therefore  $z(t) = u(t) + v(t)$ .

**Remark:** As found in [3], for  $G$  defined in (5), when  $|a| < \Gamma(\alpha - \beta + 1)$ ,  $G(0, \cdot) = G(1, \cdot) = G(\cdot, 0) = G(\cdot, 1) = 0$ , and  $|G(t, s)| \leq \overline{G}(s)$  on  $[0, 1] \times [0, 1]$  where

$$\overline{G}(s) = \begin{cases} (1-s)^{\alpha-1}E_{\alpha-\beta,\alpha}[a(1-s)^{\alpha-\beta}], & a \in [0, \Gamma(\alpha - \beta + 1)), \\ \frac{(1-s)^{\alpha-1}E_{\alpha-\beta,\alpha}[a(1-s)^{\alpha-\beta}]}{\Gamma(\alpha)E_{\alpha-\beta,\alpha}[a]}, & a \in (-\Gamma(\alpha - \beta + 1), 0). \end{cases} \quad (6)$$

We need to define the constant

$$U = \int_0^1 \overline{G}(s)w(s)ds \quad (7)$$

with  $\overline{G}$  defined in (6). We take  $X = C[0, 1]$  with the norm  $\|u\| = \max_{t \in [0, 1]} |u(t)|$ ,  $u \in X$ , to be our Banach space.

We have developed two main theorems on the existence and uniqueness of solutions for our problem.

**Theorem 1:** Let  $|a| < \Gamma(\alpha - \beta + 1)$ . Assume there exists  $r > 0$  such that

$$|f(t, x)| \leq \frac{r - k}{U} \quad \text{for } (t, x) \in [0, 1] \times [-r, r], \quad (8)$$

where  $k$  is the constant given in the right hand boundary condition given in (2). Then, BVP (1)-(2) has at least one solution  $z$  with  $\|z\| \leq r$ .

**Theorem 2:** Let  $|a| < \Gamma(\alpha - \beta + 1)$ . Assume that  $f$  satisfies the Lipschitz

condition in  $x$

$$|f(t, x_1) - f(t, x_2)| \leq B|x_1 - x_2| \quad \text{for } (t, x_1), (t, x_2) \in [0, 1] \times \mathbb{R},$$

with  $B \in (0, 1/U)$ . Then, BVP (1)-(2) has a unique solution.

### Proofs:

For the purpose of these proofs, we need to define an operator  $T : X \rightarrow X$  by

$$(Tz)(t) = \int_0^1 G(t, s)w(s)f(s, u(s))ds + \frac{kt^{\alpha-1}E_{\alpha-\beta, \alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta, \alpha}(a)},$$

where  $G$  is the Green's function used to solve for  $u(t)$ . It is clear that  $z$  is a solution of BVP (1)-(2) if and only if  $z \in X$  is a fixed point of  $T$ . There is a standard argument to show that  $T$  is completely continuous.

**Proof of Theorem 1:** Let  $\Omega \subset X$  be the set defined by

$$\Omega = \{z \in X \mid \|z\| \leq r\},$$

where  $r$  is given in Theorem 1.

For any  $z \in \Omega$ , we have  $\|z(t)\| \leq r$  on  $[0, 1]$ , and so

$$\begin{aligned} |(Tz)(t)| &= \left| \int_0^1 G(t, s)w(s)f(s, u(s))ds + \frac{kt^{\alpha-1}E_{\alpha-\beta, \alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta, \alpha}(a)} \right| \\ &\leq \left| \int_0^1 G(t, s)w(s)f(s, u(s))ds \right| + \left| \frac{kt^{\alpha-1}E_{\alpha-\beta, \alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta, \alpha}(a)} \right| \\ &\leq \int_0^1 |G(t, s)|w(s)|f(s, u(s))|ds + k \\ &\leq \int_0^1 \frac{\overline{G}(s)w(s)(r-k)}{U}ds + k = \frac{r-k}{U}U + k \\ &= r - k + k = r, \quad t \in [0, 1]. \end{aligned}$$

Hence,  $\|Tu\| \leq r$ . Therefore,  $T\Omega \subset \Omega$ .

By Schauder fixed point theorem,  $T$  has a fixed point  $z$  in  $\Omega$ . Hence, our BVP has at least one solution  $z(t)$  with  $\|z\| \leq r$ .  $\square$

**Proof of Theorem 2:** Let  $z_1, z_2 \in X$  be solutions of (0.1). Then  $z_1 = u_1 + v_1$  and  $z_2 = u_2 + v_2$ , where  $u_1, u_2$  are solutions of (0.3) and  $v_1, v_2$  are solutions of (0.4). For  $t \in [0, 1]$ ,

$$\begin{aligned}
|(Tz_1 - Tz_2)(t)| &= \left| \int_0^1 G(t, s)w(s)f(s, u_1(s))ds + \frac{kt^{\alpha-1}E_{\alpha-\beta, \alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta, \alpha}(a)} \right. \\
&\quad \left. - \int_0^1 G(t, s)w(s)f(s, u_2(s))ds - \frac{kt^{\alpha-1}E_{\alpha-\beta, \alpha}(at^{\alpha-\beta})}{E_{\alpha-\beta, \alpha}(a)} \right| \\
&= \left| \int_0^1 G(t, s)w(s)f(s, u_1(s))ds - \int_0^1 G(t, s)w(s)f(s, u_2(s))ds \right| \\
&= \left| \int_0^1 G(t, s)w(s)(f(s, u_1(s)) - f(s, u_2(s))) ds \right| \\
&\leq \int_0^1 \overline{G}(s)w(s) |f(s, u_1(s)) - f(s, u_2(s))| ds \\
&\leq \int_0^1 \overline{G}(s)w(s)B |u_1(s) - u_2(s)| ds \leq BU\|u_1 - u_2\|.
\end{aligned}$$


Since  $BU < 1$ ,  $T$  is a contraction mapping. By the contraction mapping principle,  $T$  has a unique fixed point. Therefore, our BVP has a unique solution.  $\square$

### Examples:

**Example 1:** Consider the BVP

$$\begin{cases} -D_{0+}^{3.6}z + aD_{0+}^{1.4}z = [\cos(20t) + 1 + e^{\sin(100t)}] \left[ \frac{1}{2} \sin(z) + \frac{z}{z+1} + e^t \right] \\ z(0) = z'(0) = \dots = z^{(n-2)}(0) = 0, \quad z(1) = 0.01 \end{cases} \quad (9)$$

Here,  $f(t, z) = \frac{1}{2} \sin(z) + \frac{z}{z+1} + e^t$  and  $w(t) = \cos(20t) + 1 + e^{\sin(100t)}$ . Furthermore,  $\Gamma(3.2) = 2.424$ . The following figure represents the solution to this problem using an iterative process with  $a = 2.4$  shown in green,  $a = 0.4$  shown in red, and  $a = -2.4$  shown in blue.



**Example 2:** Consider the BVP

$$\begin{cases} -D_{0+}^{4.5}z + 1.32D_{0+}^{2.7}z = \left[ \frac{z^2}{z^2+1} \right] [\cos(t) + 1] \\ z(0) = z'(0) = \dots = z^{(n-2)} = 0, \quad z(1) = 10^{-13} \end{cases} \quad (10)$$

Here,  $f(t, z) = \frac{z^2}{z^2+1}$  and  $w(t) = \cos(t)+1$ . Note that  $|a| = 1.32 < 1.6765 \approx \Gamma(2.8)$ .

We claim that this problem has a unique solution. By (0.6) and (0.7),  $1/U \approx 24.5834$ .


Then, for  $(t, z_1), (t, z_2) \in [0, 1] \times \mathbb{R}$ :

$$\begin{aligned} |f(t, z_1) - f(t, z_2)| &= \left| \frac{z_1^2}{z_1^2+1} - \frac{z_2^2}{z_2^2+1} \right| \\ &= \left| \frac{z_1^2(z_2^2+1) - z_2^2(z_1^2+1)}{(z_1^2+1)(z_2^2+1)} \right| \\ &= \left| \frac{z_1^2 - z_2^2}{(z_1^2+1)(z_2^2+1)} \right| \\ &= \left| \frac{(z_1 - z_2)(z_1 + z_2)}{(z_1^2+1)(z_2^2+1)} \right| \\ &= |z_1 - z_2| \left| \frac{z_1 + z_2}{(z_1^2+1)(z_2^2+1)} \right| \end{aligned}$$



$$\begin{aligned}
&= |z_1 - z_2| \left| \frac{z_1}{(z_1^2 + 1)(z_2^2 + 1)} + \frac{z_2}{(z_1^2 + 1)(z_2^2 + 1)} \right| \\
&< |z_1 - z_2| \left| \frac{z_1}{z_1^2 + 1} + \frac{z_2}{z_2^2 + 1} \right| \\
&< |z_1 - z_2| |1 + 1| \\
&= 2|z_1 - z_2|.
\end{aligned}$$

Therefore,  $f$  satisfies the Lipschitz condition with  $B = 2 \in (0, 1/U)$ . Hence by Theorem 2, BVP (0.10) has a unique solution. The numerical solution is given below.



## References

- [AOS] [1] R. Agarwal, D. O'Regan, and S. Staněk, Positive solutions for Dirichlet problems of singular nonlinear fractional differential equations, *J. Math. Anal. Appl.* **371** (2010), 57–68.
- [AN] [2] B. Ahmad and J. J. Nieto, Existence results for a coupled system of nonlinear fractional differential equations with three-point boundary conditions, *Comput. Math. Appl.* **58** (2009), 1838–1843.
- [AWY] [3] M. Ackerman, M. Wang, and G. Yost, A Fractional Boundary Value Problem with Separated Boundary Conditions, *Panamerican Mathematical Journal* **25** (2015), 1–12.
- [B] [4] C. Bai, Infinitely many solutions for a perturbed nonlinear fractional boundary-value problem. *Electron. J. Differential Equations* **2013**, No. 136, 12 pp.
- [BL] [5] Z. Bai and H. Lü, Positive solutions for boundary value problem of nonlinear fractional differential equation, *J. Math. Anal. Appl.* **311** (2005), 495–505.
- [CH] [6] A. Cabada and Z. Hamdi, Nonlinear fractional differential equations with integral boundary value conditions, *Appl. Math. Comput.* **228** (2014), 251–257.
- [D] [7] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, New York, 1985.
- [FZG] [8] M. Feng, X. Zhang, and W. Ge, New existence results for higher-order nonlinear fractional differential equation with integral boundary conditions, *Bound. Value Probl.* (2011), Art. ID 720702, 20 pp.
- [G] [9] C. Goodrich, Existence of a positive solution to a class of fractional differential equations, *Appl. Math. Lett.* **23** (2010), 1050–1055.

- GKKW2** [10] J. R. Graef, L. Kong, Q. Kong, and M. Wang, Uniqueness of positive solutions of fractional boundary value problems with non-homogeneous integral boundary conditions, *Fract. Calc. Appl. Anal.* **15** (2012), 509-528.
- GKKW** [11] J. R. Graef, L. Kong, Q. Kong, and M. Wang, Fractional boundary value problems with integral boundary conditions, *Appl. Anal.* **92** (2013), 2008–2020.
- GKKW3** [12] J. R. Graef, L. Kong, Q. Kong, and M. Wang, Existence and uniqueness of solutions for a fractional boundary value problem with Dirichlet boundary condition, *Electron. J. Qual. Theory Differ. Equ.* **2013** No. 55, 11 pp.
- GKKW4** [13] J. R. Graef, L. Kong, Q. Kong, and M. Wang, A fractional boundary value problem with Dirichlet boundary condition, *Commun. Appl. Anal.*, accepted.
- GKW** [14] J. R. Graef, L. Kong, and M. Wang, A Chebyshev spectral method for solving Riemann-Liouville fractional boundary value problems, *Appl. Math. Comput.* **241** (2014), 140 - 150.
- GL** [15] D. Guo and V. Lakshmikantham, *Nonlinear Problems in Abstract Cones*, Academic Press, New York, 1988.
- HL** [16] J. Henderson and R. Luca, Positive solutions for a system of nonlocal fractional boundary value problems, *Fract. Calc. Appl. Anal.* **16** (2013), 985–1008.
- H** [17] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific, Singapore, 2000.
- JY** [18] D. Jiang and C. Yuan, The positive properties of the Green function for Dirichlet-type boundary value problems of nonlinear fractional differential equations and its application, *Nonlinear Anal.* **72** (2010), 710–719.
- KST** [19] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, Elsevier, 2006.

- [KKW] [20] L. Kong, Q. Kong, and M. Wang, Existence and uniqueness of solutions for a fractional boundary value problem with a separated boundary condition, *Dynamic Systems and Applications*, **23** (2014), 691–698.
- [KW0] [21] Q. Kong and M. Wang, Positive solutions of nonlinear fractional boundary value problems with Dirichlet boundary conditions, *J. Qual. Theory Differ. Equ.* **No. 17** (2012), 1–13.
- [Ta] [22] V. Tarasov, *Fractional Dynamics: Applications of Fractional Calculus to Dynamics of Particles, Fields and Media*, Springer-Verlag, New York, 2011.
- [YC] [23] L. Yang and H. Chen, Unique positive solutions for fractional differential equation boundary value problems, *Appl. Math. Lett.* **23** (2010), 1095–1098.
- [Z] [24] E. Zeidler, *Nonlinear Functional Analysis and its Applications I: Fixed-Point Theorems*, Springer-Verlag, New York, 1986.
- [Zh] [25] S. Zhang, Positive solutions to singular boundary value problem for nonlinear fractional differential equation, *Comput. Math. Appl.* **59** (2010), 1300–1309.